

PATIENT INFORMATION
OHTUVAYRE (OH-too-vare)
(ensifentrine)
inhalation suspension, for oral inhalation

Important: For oral inhalation only. Do not swallow or inject OHTUVAYRE.

What is OHTUVAYRE?

- OHTUVAYRE is a prescription medicine used to treat chronic obstructive pulmonary disease (COPD) in adults. COPD is a chronic (long-term) lung disease that includes chronic bronchitis, emphysema, or both.
- OHTUVAYRE is an inhibitor of phosphodiesterases 3 and 4 (PDE3 and PDE4).
- Decreasing PDE3 activity helps the muscles around the airways in your lungs stay relaxed to prevent symptoms such as wheezing, coughing, chest tightness, and shortness of breath, which can happen when muscles around the airway tighten, making it hard to breathe. Decreasing PDE4 activity helps to decrease inflammation in the lungs. Inflammation in the lungs can lead to breathing problems.
- OHTUVAYRE is used to improve symptoms of COPD for better breathing and to reduce the number of flare-ups (the worsening of your COPD symptoms for several days).
- OHTUVAYRE is used long-term as 1 ampule of OHTUVAYRE, 2 times each day (1 in the morning and 1 in the evening) inhaled through your nebulizer that is fitted with a mouthpiece.
- **OHTUVAYRE is not used to relieve sudden breathing problems and will not replace an inhaled rescue medicine.**
- **OHTUVAYRE should not be used in children.** It is not known if OHTUVAYRE is safe and effective in children.

Do not use OHTUVAYRE if you have had an allergic reaction to ensifentrine or any of the ingredients in OHTUVAYRE. See “**What are the ingredients in OHTUVAYRE?**” at the end of this Patient Information leaflet for a complete list of ingredients in OHTUVAYRE.

Before taking OHTUVAYRE, tell your healthcare provider about all your medical conditions, including if you:

- have or have had a history of mental health problems including depression and suicidal behavior.
- have liver problems.
- are pregnant or plan to become pregnant. It is not known if OHTUVAYRE may harm your unborn baby.
- are breastfeeding. It is not known if the medicine in OHTUVAYRE passes into your breast milk and if it can harm your baby.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How should I use OHTUVAYRE?

Read the step-by-step instructions for using OHTUVAYRE at the end of this Patient Information leaflet.

- **OHTUVAYRE is only for use with a nebulizer.**
- **Do not** use OHTUVAYRE unless your healthcare provider has taught you how to use it with your nebulizer and you understand how to use it correctly.
- Use OHTUVAYRE exactly as your healthcare provider tells you to use it. **Do not** use OHTUVAYRE more often than prescribed.
- OHTUVAYRE is taken as a breathing treatment by oral (mouth) inhalation and should be used with a standard jet nebulizer with a mouthpiece connected to an air compressor.
- **Do not** mix other medicines with OHTUVAYRE in your nebulizer.

- **Shake the OHTUVAYRE ampule vigorously immediately before using it. When shaken, OHTUVAYRE will appear cloudy and yellow to pale yellow in color.**
- Use 1 ampule of OHTUVAYRE 2 times a day, 1 time in the morning and 1 time in the evening. **Do not** use more than 1 ampule of OHTUVAYRE in a single treatment. **Do not** use more than 2 ampules of OHTUVAYRE a day.
- If you use too much OHTUVAYRE, call your healthcare provider or go to the nearest hospital emergency room right away if you have any unusual symptoms, such as increased heart rate or headaches.
- **OHTUVAYRE does not relieve sudden symptoms of COPD and you should not use extra doses of OHTUVAYRE to relieve these sudden symptoms.** Always have an inhaled rescue medicine with you to treat sudden symptoms. If you do not have an inhaled rescue medicine, call your healthcare provider to have one prescribed for you.
- Call your healthcare provider or get emergency medical care right away if:
 - your breathing problems get worse.
 - you need to use your inhaled rescue medicine more often than usual.
 - your inhaled rescue medicine does not relieve your symptoms.
- **Do not** stop using OHTUVAYRE, even if you are feeling better, unless your healthcare provider tells you to, because your symptoms might get worse.
- Throw away the OHTUVAYRE ampules immediately after use. Due to their small size, the ampules pose a danger of choking to young children.

What are the possible side effects of OHTUVAYRE?

OHTUVAYRE can cause serious side effects, including:

- **Sudden breathing problems immediately after inhaling your medicine.** If you have sudden breathing problems immediately after inhaling your medicine, stop using OHTUVAYRE and call your healthcare provider right away or go to the nearest hospital emergency room right away.
- **Mental health problems including suicidal thoughts and behavior.** You may experience mood or behavior changes when taking OHTUVAYRE. Call your healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:
 - thoughts of suicide or dying
 - attempt to commit suicide
 - trouble sleeping (insomnia)
 - new or worse anxiety
 - new or worse depression
 - acting on dangerous impulses
 - other unusual changes in your behavior or mood

Most common side effects of OHTUVAYRE include:

- back pain
- high blood pressure
- bladder infection
- diarrhea

These are not all the possible side effects of OHTUVAYRE.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store OHTUVAYRE?

- Store OHTUVAYRE at room temperature between 68°F to 77°F (20°C to 25°C). Keep OHTUVAYRE away from direct sunlight and excessive heat. Do not freeze.
- Store OHTUVAYRE in the unopened protective foil pouch and **only open the foil pouch immediately before using OHTUVAYRE.** For pouches of 5 ampules, remove one ampule and place the remaining ampules back into the pouch until next use.

- Throw away the used ampule and any leftover medicine after use.
- Do not use OHTUVAYRE after the expiration date provided on the foil pouch and ampule.
- **Keep OHTUVAYRE and all medicines out of the reach of children.**

General information about the safe and effective use of OHTUVAYRE.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use OHTUVAYRE for a condition for which it was not prescribed. Do not give OHTUVAYRE to other people, even if they have the same symptoms that you have. It may harm them.

You can ask your healthcare provider or pharmacist for information about OHTUVAYRE that is written for health professionals.

What are the ingredients in OHTUVAYRE?

Active ingredient: ensifentrine

Inactive ingredients: dibasic sodium phosphate, monobasic sodium phosphate, polysorbate 20, sodium chloride, sorbitan monolaurate and water for injection.

Manufactured for:
Verona Pharma, Inc.
Raleigh, NC, 27615

©2025. Verona Pharma plc and its affiliates

Verona Pharma, OHTUVAYRE and the Verona Pharma and OHTUVAYRE logos are all trademarks of Verona Pharma plc

This Patient Information has been approved by the U.S. Food and Drug Administration. Approved: 7/2025

PIL002 REF-00414